

# DYNAMICS OF TUPLES OF MATRICES

G. COSTAKIS, D. HADJILOUCAS, AND A. MANOUSSOS

**ABSTRACT.** In this article we answer a question raised by N. Feldman in [4] concerning the dynamics of tuples of operators on  $\mathbb{R}^n$ . In particular, we prove that for every positive integer  $n \geq 2$  there exist  $n$  tuples  $(A_1, A_2, \dots, A_n)$  of  $n \times n$  matrices over  $\mathbb{R}$  such that  $(A_1, A_2, \dots, A_n)$  is hypercyclic. We also establish related results for tuples of  $2 \times 2$  matrices over  $\mathbb{R}$  or  $\mathbb{C}$  being in Jordan form.

## 1. INTRODUCTION

Following the recent work of Feldman in [4], an  $n$ -tuple of operators is a finite sequence of length  $n$  of commuting continuous linear operators  $T_1, T_2, \dots, T_n$  acting on a locally convex space  $X$ . The tuple  $(T_1, T_2, \dots, T_n)$  is hypercyclic if there exists a vector  $x \in X$  such that the set

$$\{T_1^{k_1} T_2^{k_2} \dots T_n^{k_n} x : k_1, k_2, \dots, k_n \geq 0\}$$

is dense in  $X$ . Such a vector  $x$  is called hypercyclic for  $(T_1, T_2, \dots, T_n)$  and the set of hypercyclic vectors for  $(T_1, T_2, \dots, T_n)$  will be denoted by  $HC((T_1, T_2, \dots, T_n))$ . The above definition generalizes the notion of hypercyclicity to tuples of operators. For an account of results, comments and an extensive bibliography on hypercyclicity we refer to [1], [5], [6] and [7]. For results concerning the dynamics of tuples of operators see [2], [3], [4] and [8].

In [4] Feldman showed, among other things, that in  $\mathbb{C}^n$  there exist diagonalizable  $n+1$ -tuples of matrices having dense orbits. In addition he proved that there is no  $n$ -tuple of diagonalizable matrices on  $\mathbb{R}^n$  or  $\mathbb{C}^n$  that has a somewhere dense orbit. Therefore the following question arose naturally.

**Question (Feldman [4]).** *Are there non-diagonalizable  $n$ -tuples on  $\mathbb{R}^k$  that have somewhere dense orbits?*

---

2000 *Mathematics Subject Classification.* 47A16.

*Key words and phrases.* Hypercyclic operators, tuples of matrices.

During this research the third author was fully supported by SFB 701 "Spektrale Strukturen und Topologische Methoden in der Mathematik" at the University of Bielefeld, Germany. He would also like to express his gratitude to Professor H. Abels for his support.

We give a positive answer to this question in a very strong form, as the next theorem shows.

**Theorem 1.1.** *For every positive integer  $n \geq 2$  there exist  $n$ -tuples  $(A_1, A_2, \dots, A_n)$  of  $n \times n$  non-diagonalizable matrices over  $\mathbb{R}$  such that  $(A_1, A_2, \dots, A_n)$  is hypercyclic.*

Restricting ourselves to tuples of  $2 \times 2$  matrices in Jordan form either on  $\mathbb{R}^2$  or  $\mathbb{C}^2$ , we prove the following.

**Theorem 1.2.** *There exist  $2 \times 2$  matrices  $A_j, j = 1, 2, 3, 4$  in Jordan form over  $\mathbb{R}$  such that  $(A_1, A_2, A_3, A_4)$  is hypercyclic. In particular*

$$HC((A_1, A_2, A_3, A_4)) = \left\{ \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2 : x_2 \neq 0 \right\}.$$

**Theorem 1.3.** *There exist  $2 \times 2$  matrices  $A_j, j = 1, 2, \dots, 8$  in Jordan form over  $\mathbb{C}$  such that  $(A_1, A_2, \dots, A_8)$  is hypercyclic.*

## 2. PRODUCTS OF $2 \times 2$ MATRICES

**Lemma 2.1.** *Let  $m$  be a positive integer and for each  $j = 1, 2, \dots, m$  let  $A_j$  be a  $2 \times 2$  matrix in Jordan form over a field  $\mathbb{F} = \mathbb{C}$  or  $\mathbb{R}$ , i.e.*

*$A_j = \begin{pmatrix} a_j & 1 \\ 0 & a_j \end{pmatrix}$  for  $a_1, a_2, \dots, a_m \in \mathbb{F}$ . Then  $(A_1, A_2, \dots, A_m)$  over  $\mathbb{C}$  (respectively  $\mathbb{R}$ ) is hypercyclic if and only if the sequence*

$$\left\{ \begin{pmatrix} \frac{k_1}{a_1} + \frac{k_2}{a_2} + \dots + \frac{k_m}{a_m} \\ a_1^{k_1} a_2^{k_2} \dots a_m^{k_m} \end{pmatrix} : k_1, k_2, \dots, k_m \in \mathbb{N} \right\}$$

*is dense in  $\mathbb{C}^2$  (respectively  $\mathbb{R}^2$ ).*

*Proof.* We prove the above in the case  $\mathbb{F} = \mathbb{C}$ , since the other case is similar. Observe that

$$A_j^l = \begin{pmatrix} a_j^l & l a_j^{l-1} \\ 0 & a_j^l \end{pmatrix}$$

for  $l \in \mathbb{N}$ . As a result we have

$$A_1^{k_1} A_2^{k_2} \dots A_m^{k_m} = \begin{pmatrix} \prod_{j=1}^m a_j^{k_j} & \prod_{j=1}^m a_j^{k_j} \sum_{s=1}^m \frac{k_s}{a_s} \\ 0 & \prod_{j=1}^m a_j^{k_j} \end{pmatrix}.$$

Assume that  $(A_1, A_2, \dots, A_m)$  is hypercyclic and let  $\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \in \mathbb{C}^2$  be a hypercyclic vector for  $(A_1, A_2, \dots, A_m)$ . Then the sequence

$$\left\{ A_1^{k_1} A_2^{k_2} \dots A_m^{k_m} \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} : k_1, k_2, \dots, k_m \in \mathbb{N} \right\}$$

$$= \left\{ \begin{pmatrix} z_1 \prod_{j=1}^m a_j^{k_j} + z_2 \prod_{j=1}^m a_j^{k_j} \sum_{s=1}^m \frac{k_s}{a_s} \\ z_2 \prod_{j=1}^m a_j^{k_j} \end{pmatrix} : k_1, k_2, \dots, k_m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{C}^2$ . This implies that  $z_2 \neq 0$ . Dividing the element on the first row by that on the second, it can easily be shown that the sequence

$$\left\{ \begin{pmatrix} \frac{k_1}{a_1} + \frac{k_2}{a_2} + \dots + \frac{k_m}{a_m} \\ a_1^{k_1} a_2^{k_2} \dots a_m^{k_m} \end{pmatrix} : k_1, k_2, \dots, k_m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{C}^2$ . The converse can easily be shown.  $\square$

*Remark 2.2.* Let  $m$  be a positive integer and for each  $j = 1, 2, \dots, m$  let  $A_j$  be a  $2 \times 2$  matrix in Jordan form over a field  $\mathbb{F} = \mathbb{C}$  or  $\mathbb{R}$ . By the proof of Lemma 2.1 it is immediate that whenever  $(A_1, A_2, \dots, A_m)$  over  $\mathbb{C}$  (respectively  $\mathbb{R}$ ) is hypercyclic one can completely describe the set of hypercyclic vectors as

$$\left\{ \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \in \mathbb{C}^2 : z_2 \neq 0 \right\} \quad \left( \text{respectively } \left\{ \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2 : x_2 \neq 0 \right\} \right).$$

**2.1. The real case.** We shall need the following well known result, see for example [4].

**Theorem 2.3.** *If  $a, b > 1$  and  $\frac{\ln a}{\ln b}$  is irrational then the sequence  $\{\frac{a^n}{b^m} : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{R}^+$ .*

**Lemma 2.4.** *Let  $a, b \in \mathbb{R}$  such that  $-1 < a < 0$ ,  $b > 1$  and  $\frac{\ln |a|}{\ln b}$  is irrational. Then the sequence  $\{a^n b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{R}$ .*

*Proof.* Since  $\frac{\ln |a|}{\ln b}$  is irrational it follows that  $\ln b / \ln \frac{1}{a^2}$  is irrational as well. Applying Theorem 2.3 we conclude that the sequence  $\{a^{2n} b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{R}^+$ . On the other hand the fact that  $a$  is negative implies that the sequence  $\{a^{2n+1} b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{R}^-$ . This completes the proof of the lemma.  $\square$

In what follows, for any  $x \in \mathbb{R}$  we will be denoting by  $[x]$  the ‘integral part of  $x$ ’, that is, the largest integer which does not exceed  $x$ , and by  $\{x\}$  the ‘fractional part of  $x$ ’, that is,  $\{x\} = x - [x]$ .

**Proposition 2.5.** *There exist  $a_1, a_2, a_3, a_4 \in \mathbb{R}$  such that the sequence*

$$\left\{ \begin{pmatrix} \frac{k_1}{a_1} + \frac{k_2}{a_2} + \frac{k_3}{a_3} + \frac{k_4}{a_4} \\ a_1^{k_1} a_2^{k_2} a_3^{k_3} a_4^{k_4} \end{pmatrix} : k_1, k_2, k_3, k_4 \in \mathbb{N} \right\}$$

*is dense in  $\mathbb{R}^2$ .*

*Proof.* By the lemma above fix  $a, b \in \mathbb{R}$  such that  $-1 < a < 0$ ,  $a + \frac{1}{a} \in \mathbb{R} \setminus \mathbb{Q}$  and  $\{a^n b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{R}$ . Let  $x_1, x_2 \in \mathbb{R}$  and  $\epsilon > 0$  be given. Then there exist  $n, m \in \mathbb{N}$  such that  $|a^n b^m - x_2| < \epsilon$ . Note that  $a^n b^m = a^{n+k} b^m \frac{1}{a^k} 1^s$  for every  $k, s \in \mathbb{N}$ . Note also that  $a + \frac{1}{a} < 0$ . In the case  $x_1 \geq 0$  (resp.  $x_1 < 0$ ) there exists  $k \in \mathbb{N}$  such that  $\frac{n}{a} + \frac{m}{b} + k \left(a + \frac{1}{a}\right)$  is less than  $-1$  (resp.  $x_1 - 1$ ) and

$$\left| \left\{ \frac{n}{a} + \frac{m}{b} + k \left(a + \frac{1}{a}\right) \right\} - \{x_1\} \right| < \epsilon.$$

As there exists  $s \in \mathbb{N}$  such that

$$\left| \frac{n}{a} + \frac{m}{b} + k \left(a + \frac{1}{a}\right) + s - x_1 \right| < \epsilon$$

we are done. Hence, setting  $a_1 = a, a_2 = b, a_3 = \frac{1}{a}, a_4 = 1$  the result is proved.  $\square$

*Proof of Theorem 1.2.* This is an immediate consequence of Lemma 2.1, Proposition 2.5 and Remark 2.2.

**Example 2.6.** One may construct many concrete examples of four  $2 \times 2$  matrices, in Jordan form over  $\mathbb{R}$ , being hypercyclic. For example, fix  $a, b \in \mathbb{R}$  such that  $-1 < a < 0$ ,  $b > 1$  and both  $a + \frac{1}{a}$ ,  $\frac{\ln|a|}{\ln b}$  are irrational. From the above we conclude that

$$\left( \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}, \begin{pmatrix} b & 1 \\ 0 & b \end{pmatrix}, \begin{pmatrix} \frac{1}{a} & 1 \\ 0 & \frac{1}{a} \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right)$$

is hypercyclic.

**Proposition 2.7.** (i) Every pair  $(A_1, A_2)$  of  $2 \times 2$  matrices over  $\mathbb{R}$  with  $A_j$ ,  $j = 1, 2$  being either diagonal or in Jordan form is not hypercyclic.

(ii) There exist pairs  $(A_1, A_2)$  of  $2 \times 2$  matrices over  $\mathbb{R}$  such that  $A_1$  is diagonal,  $A_2$  is antisymmetric (rotation matrix) and  $(A_1, A_2)$  is hypercyclic. In particular every non-zero vector in  $\mathbb{R}^2$  is hypercyclic for  $(A_1, A_2)$ , i.e.

$$HC((A_1, A_2)) = \mathbb{R}^2 \setminus \{(0, 0)\}.$$

(iii) There exist pairs  $(A_1, A_2)$  of  $2 \times 2$  matrices over  $\mathbb{R}$  such that both  $A_1$  and  $A_2$  are antisymmetric and  $(A_1, A_2)$  is hypercyclic. In particular every non-zero vector in  $\mathbb{R}^2$  is hypercyclic for  $(A_1, A_2)$ , i.e.

$$HC((A_1, A_2)) = \mathbb{R}^2 \setminus \{(0, 0)\}.$$

*Proof.* Let us prove assertion (i). The case of  $A_1, A_2$  both diagonal is covered by Feldman, see [4]. Assume that  $A_1$  is diagonal and  $A_2$  is in Jordan form, i.e.  $A_1 = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ ,  $A_2 = \begin{pmatrix} b & 1 \\ 0 & b \end{pmatrix}$  for  $a, b \in$

$\mathbb{R}$ . Suppose that  $(A_1, A_2)$  is hypercyclic and let  $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \in \mathbb{R}^2$  be a hypercyclic vector for  $(A_1, A_2)$ . Then the sequence

$$\left\{ A_1^n A_2^m \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} : n, m \in \mathbb{N} \right\},$$

i.e. the sequence  $\left\{ \begin{pmatrix} a^n b^m x_1 + m a^n b^m x_2 \\ a^n b^m x_2 \end{pmatrix} : n, m \in \mathbb{N} \right\}$  is dense in  $\mathbb{R}^2$ . Observe that  $x_2$  cannot be zero. Then there exist  $y_1, y_2 \in \mathbb{R}$  and sequences of positive integers  $\{n_k\}, \{m_k\}$  such that  $m_k \rightarrow +\infty$  and

$$a^{n_k} b^{m_k} x_1 + m_k a^{n_k} b^{m_k} x_2 \rightarrow y_1,$$

$$a^{n_k} b^{m_k} x_2 \rightarrow y_2$$

as  $k \rightarrow +\infty$ . It clearly follows that  $|m_k a^{n_k} b^{m_k} x_2| \rightarrow +\infty$  which is a contradiction. Assume now that both  $A_1, A_2$  are in Jordan form, i.e.

$$A_1 = \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}, \quad A_2 = \begin{pmatrix} b & 1 \\ 0 & b \end{pmatrix},$$

for  $a, b \in \mathbb{R}$  and  $(A_1, A_2)$  is hypercyclic. Lemma 2.1 implies that the sequence

$$\left\{ \begin{pmatrix} \frac{n}{a} + \frac{m}{b} \\ a^n b^m \end{pmatrix} : n, m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{R}^2$ . Observe that both  $|a|, |b|$  are not equal to 1. By taking absolute value in the second coordinate and then applying the logarithmic function, it follows that the sequence

$$\left\{ \begin{pmatrix} \frac{n}{a} + \frac{m}{b} \\ n \ln |a| + m \ln |b| \end{pmatrix} : n, m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{R}^2$ . Hence the sequence

$$\left\{ \begin{pmatrix} n \frac{\ln |a|}{a} + m \frac{\ln |a|}{b} \\ n \frac{\ln |a|}{a} + m \frac{\ln |b|}{a} \end{pmatrix} : n, m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{R}^2$ . Subtracting the second coordinate from the first one we conclude that the sequence

$$\left\{ m \left( \frac{\ln |a|}{b} - \frac{\ln |b|}{a} \right) : m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{R}$  which is absurd. We proceed with the proof of assertion (ii). There exist  $a \in \mathbb{R} \setminus \mathbb{Q}$  and  $b \in \mathbb{C}$  such that the sequence  $\{a^n b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{C}$ , see [4]. Write  $b = |b|e^{i\theta}$  and set

$$A_1 = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}, \quad A_2 = \begin{pmatrix} |b| \cos \theta & -|b| \sin \theta \\ |b| \sin \theta & |b| \cos \theta \end{pmatrix}.$$

Then we have

$$A_1^n A_2^m = \begin{pmatrix} a^n |b|^m \cos(m\theta) & -a^n |b|^m \sin(m\theta) \\ a^n |b|^m \sin(m\theta) & a^n |b|^m \cos(m\theta) \end{pmatrix}.$$

Applying in the above relation the vector  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and taking into account that the sequence  $\{a^n b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{C}$  we conclude that the sequence

$$\left\{ A_1^n A_2^m \begin{pmatrix} 1 \\ 0 \end{pmatrix} : n, m \in \mathbb{N} \right\} = \left\{ \begin{pmatrix} a^n |b|^m \cos(m\theta) \\ a^n |b|^m \sin(m\theta) \end{pmatrix} : n, m \in \mathbb{N} \right\}$$

is dense in  $\mathbb{R}^2$ . Hence  $(A_1, A_2)$  is hypercyclic. It is now easy to show that every non-zero vector in  $\mathbb{R}^2$  is hypercyclic for  $(A_1, A_2)$ .

In order to prove the last assertion we follow a similar line of reasoning as above. That is, by Corollary 3.2 in [4] there exist  $a, b \in \mathbb{C} \setminus \mathbb{R}$  such that the sequence  $\{a^n b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{C}$ . Write  $a = |a|e^{i\phi}$ ,  $b = |b|e^{i\theta}$  and set

$$A_1 = \begin{pmatrix} |a| \cos \phi & -|a| \sin \phi \\ |a| \sin \phi & |a| \cos \phi \end{pmatrix}, \quad A_2 = \begin{pmatrix} |b| \cos \theta & -|b| \sin \theta \\ |b| \sin \theta & |b| \cos \theta \end{pmatrix}.$$

A direct computation gives that  $\left\{ A_1^n A_2^m \begin{pmatrix} 1 \\ 0 \end{pmatrix} : n, m \in \mathbb{N} \right\}$  equals to  $\left\{ \begin{pmatrix} |a|^n |b|^m \cos(n\phi - m\theta) \\ |a|^n |b|^m \sin(n\phi + m\theta) \end{pmatrix} : n, m \in \mathbb{N} \right\}$  and by the choice of  $a, b$  we conclude that the vector  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  is hypercyclic for  $(A_1, A_2)$ . This completes the proof of the proposition.  $\square$

**Question 2.8.** *What is the minimum number of  $2 \times 2$  matrices over  $\mathbb{R}$  in Jordan form so that their tuple form a hypercyclic operator?*

**2.2. The complex case.** In what follows we will be writing  $\Re(z)$  and  $\Im(z)$  for the real and imaginary parts of a complex number  $z$  respectively.

**Proposition 2.9.** *There exist  $a_j \in \mathbb{C}$ ,  $j = 1, 2, \dots, 8$  such that the sequence*

$$\left\{ \begin{pmatrix} \frac{k_1}{a_1} + \frac{k_2}{a_2} + \dots + \frac{k_8}{a_8} \\ a_1^{k_1} a_2^{k_2} \dots a_8^{k_8} \end{pmatrix} : k_1, k_2, \dots, k_8 \in \mathbb{N} \right\}$$

*is dense in  $\mathbb{C}^2$ .*

*Proof.* The proof is in the same spirit as the proof of Proposition 2.5. Fix  $a, b \in \mathbb{C}$  such that  $-1 < a < 0$ ,  $a + \frac{1}{a}, a - \frac{1}{a} \in \mathbb{R} \setminus \mathbb{Q}$  and  $\{a^n b^m :$

$n, m \in \mathbb{N}$  is dense in  $\mathbb{C}$  (see Corollary 3.2). Let  $z_1, z_2 \in \mathbb{C}$  and  $\epsilon > 0$  be given. Then there exist  $n, m \in \mathbb{N}$  such that  $|a^n b^m - z_2| < \epsilon$ . Note that

$$a^n b^m = a^{n+k} b^m \frac{1}{a^k} 1^s (ia)^\xi \left(\frac{1}{ia}\right)^\xi (4i)^\rho \left(-\frac{1}{4}\right)^\rho$$

for every  $k, s, \xi \in \mathbb{N}$  and  $\rho \in 4\mathbb{N}$ . Note that  $a + \frac{1}{a} < 0$  and  $a - \frac{1}{a} > 0$ . In the case  $\Re(z_1) \geq 0$  (resp.  $\Re(z_1) < 0$ ) there exists  $k \in \mathbb{N}$  such that  $\Re\left(\frac{n}{a} + \frac{m}{b} + k\left(a + \frac{1}{a}\right)\right)$  is less than  $-1$  (resp.  $\Re(z_1) - 1$ ) and

$$\left| \left\{ \Re\left(\frac{n}{a} + \frac{m}{b} + k\left(a + \frac{1}{a}\right)\right) \right\} - \{\Re(z_1)\} \right| < \epsilon.$$

Also, in the case  $\Im(z_1) \geq 0$  (resp.  $\Im(z_1) < 0$ ) there exists  $\xi \in \mathbb{N}$  such that  $\Im\left(\frac{n}{a} + \frac{m}{b} + i\xi\left(a - \frac{1}{a}\right)\right)$  is greater than  $1$  (resp.  $\Im(z_1) + 1$ ) and

$$\left| \left\{ \Im\left(\frac{n}{a} + \frac{m}{b} + i\xi\left(a - \frac{1}{a}\right)\right) \right\} - \{\Im(z_1)\} \right| < \epsilon.$$

As there exists  $\rho \in 4\mathbb{N}$  such that

$$\left| \Im\left(\frac{n}{a} + \frac{m}{b} + i\xi\left(a - \frac{1}{a}\right) - \rho\left(\frac{i}{4}\right)\right) - \Im(z_1) \right| < \epsilon$$

we are done with the imaginary part. Also, there exists  $s \in \mathbb{N}$  such that

$$\left| \Re\left(\frac{n}{a} + \frac{m}{b} + k\left(a + \frac{1}{a}\right) - 4\rho + s\right) - \Re(z_1) \right| < \epsilon.$$

But this means that the real and imaginary parts of the complex number

$$\frac{n}{a} + \frac{m}{b} + k\left(a + \frac{1}{a}\right) + s + i\xi\left(a - \frac{1}{a}\right) - \rho\frac{i}{4} - 4\rho$$

are within  $\epsilon$  of the real and imaginary parts of  $z_1$ . Hence, setting  $a_1 = a, a_2 = b, a_3 = \frac{1}{a}, a_4 = 1, a_5 = ia, a_6 = \frac{1}{ia}, a_7 = 4i, a_8 = -\frac{1}{4}$  the result is proved.  $\square$

*Proof of Theorem 1.3.* By Proposition 2.9, Lemma 2.1 and Remark 2.2 the assertion follows.

**Example 2.10.** Fix  $a, b \in \mathbb{C}$  such that  $-1 < a < 0, a + \frac{1}{a}, a - \frac{1}{a} \in \mathbb{R} \setminus \mathbb{Q}$  and  $\{a^n b^m : n, m \in \mathbb{N}\}$  is dense in  $\mathbb{C}$ . From the above it is evident that the following 8-tuple of  $2 \times 2$  matrices in Jordan form over  $\mathbb{C}$

$$\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}, \begin{pmatrix} b & 1 \\ 0 & b \end{pmatrix}, \begin{pmatrix} \frac{1}{a} & 1 \\ 0 & \frac{1}{a} \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \\ \begin{pmatrix} ia & 1 \\ 0 & ia \end{pmatrix}, \begin{pmatrix} \frac{1}{ia} & 1 \\ 0 & \frac{1}{ia} \end{pmatrix}, \begin{pmatrix} 4i & 1 \\ 0 & 4i \end{pmatrix}, \begin{pmatrix} -\frac{1}{4} & 1 \\ 0 & -\frac{1}{4} \end{pmatrix}$$

is hypercyclic.

**Question 2.11.** *What is the minimum number of  $2 \times 2$  matrices over  $\mathbb{C}$  in Jordan form so that their tuple form a hypercyclic operator?*

### 3. PRODUCTS OF $3 \times 3$ MATRICES

**Proposition 3.1. (Feldman)** *If  $b_1, b_2 \in \mathbb{D} \setminus \{0\}$  then there exists a dense set  $\Delta \subset \mathbb{C} \setminus \mathbb{D}$  such that for every  $a_1, a_2 \in \Delta$  the sequence*

$$\left\{ \begin{pmatrix} a_1^n b_1^m \\ a_2^n b_2^l \end{pmatrix} : n, m, l \in \mathbb{N} \right\}$$

*is dense in  $\mathbb{C}^2$ .*

**Corollary 3.2.** *There exist  $a \in \mathbb{C}$  and  $b, c, d \in \mathbb{R}$  such that the sequence*

$$\left\{ \begin{pmatrix} a^n b^m \\ c^n d^l \end{pmatrix} : n, m, l \in \mathbb{N} \right\}$$

*is dense in  $\mathbb{C} \times \mathbb{R}$ .*

*Proof.* Fix two real numbers  $b_1, b_2$  with  $b_1, b_2 \in (0, 1)$ . By Proposition 3.1 there exist  $a_1, a_2 \in \mathbb{C} \setminus \mathbb{D}$  such that the sequence

$$\left\{ \begin{pmatrix} a_1^n b_1^m \\ a_2^n b_2^l \end{pmatrix} : n, m, l \in \mathbb{N} \right\}$$

is dense in  $\mathbb{C}^2$ . Define  $a = a_1$ ,  $b = b_1$ ,  $c = |a_2|$  and  $d = -\sqrt{b_2}$ . Observe that the sequence

$$\left\{ \begin{pmatrix} a^n b^m \\ c^n b_2^l \end{pmatrix} : n, m, l \in \mathbb{N} \right\}$$

is dense in  $\mathbb{C} \times [0, +\infty)$ . Take  $z \in \mathbb{C}$  and  $x \in \mathbb{R}$ .

**Case I:**  $x \geq 0$ .

Then there exist sequences of positive integers  $\{n_k\}, \{m_k\}, \{l_k\}$  such that

$$a^{n_k} b^{m_k} \rightarrow z \text{ and } c^{n_k} b_2^{l_k} \rightarrow x.$$

Since  $b_2^{l_k} = d^{2l_k}$  we get  $c^{n_k} d^{2l_k} \rightarrow x$ .

**Case II:**  $x < 0$ .

Then there exist sequences of positive integers  $\{n_k\}, \{m_k\}, \{l_k\}$  such that

$$a^{n_k} b^{m_k} \rightarrow z \text{ and } c^{n_k} b_2^{l_k} \rightarrow \frac{x}{d}.$$

The last implies that  $c^{n_k} d^{2l_k+1} \rightarrow x$ . This completes the proof of the corollary.  $\square$

**Proposition 3.3.** *There exist 3 tuples  $(A_1, A_2, A_3)$  of  $3 \times 3$  matrices over  $\mathbb{R}$  such that  $(A_1, A_2, A_3)$  is hypercyclic.*



*Proof.* By Corollary 3.2 there exist  $a \in \mathbb{C}$  and  $b, c, d \in \mathbb{R}$  such that the sequence

$$\left\{ \begin{pmatrix} a^n b^m \\ c^n d^l \end{pmatrix} : n, m, l \in \mathbb{N} \right\}$$

is dense in  $\mathbb{C} \times \mathbb{R}$ . Write  $a = |a|e^{i\theta}$  and set

$$A_1 = \begin{pmatrix} |a| \cos \theta & -|a| \sin \theta & 0 \\ |a| \sin \theta & |a| \cos \theta & 0 \\ 0 & 0 & c \end{pmatrix}, \quad A_2 = \begin{pmatrix} b & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ and}$$

$$A_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & d \end{pmatrix}. \text{ Then we have}$$

$$A_1^n A_2^m A_3^l = \begin{pmatrix} |a|^n b^m \cos(n\theta) & -|a|^n b^m \sin(n\theta) & 0 \\ |a|^n b^m \sin(n\theta) & |a|^n b^m \cos(n\theta) & 0 \\ 0 & 0 & c^n d^l \end{pmatrix},$$

which in turn gives

$$A_1^n A_2^m A_3^l \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} |a|^n b^m \cos(n\theta) \\ |a|^n b^m \sin(n\theta) \\ c^n d^l \end{pmatrix}.$$

The last and the choice of  $a, b, c, d$  imply that  $(A_1, A_2, A_3)$  is hypercyclic

with  $\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$  being a hypercyclic vector for  $(A_1, A_2, A_3)$ .  $\square$

#### 4. PROOF OF THEOREM 1.1

By Proposition 2.7, there exist  $2 \times 2$  matrices  $B_1$  and  $B_2$  such that  $(B_1, B_2)$  is hypercyclic.

**Case I:**  $n = 2k$  for some positive integer  $k$ . For  $k = 1$  the result follows by Proposition 2.7. Assume that  $k > 1$ . Each  $A_j$  will be constructed by blocks of  $2 \times 2$  matrices. Let  $I_2$  be the  $2 \times 2$  identity matrix. We will be using the notation  $\text{diag}(D_1, D_2, \dots, D_n)$  to denote the diagonal matrix with diagonal entries the block matrices  $D_1, D_2, \dots, D_n$ . Define  $A_1 = \text{diag}(B_1, I_2, \dots, I_2)$ ,  $A_2 = \text{diag}(B_2, I_2, \dots, I_2)$ ,  $A_3 = \text{diag}(I_2, B_1, I_2, \dots, I_2)$ ,  $A_4 = \text{diag}(I_2, B_2, I_2, \dots, I_2)$  and so on up to  $A_{n-1} = \text{diag}(I_2, \dots, I_2, B_1)$ ,  $A_n = \text{diag}(I_2, \dots, I_2, B_2)$ . It is now easy to check that  $(A_1, A_2, \dots, A_n)$  is hypercyclic and furthermore that the set  $HC((A_1, A_2, \dots, A_n))$  is

$$\{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : x_{2j-1}^2 + x_{2j}^2 \neq 0, \forall j = 1, 2, \dots, k\}.$$

**Case II:**  $n = 2k + 1$  for some positive integer  $k$ . If  $k = 1$  the result follows by Proposition 3.3. Suppose  $k > 1$ . For simplicity we treat the case  $k = 2$ , since the general case follows by similar arguments. By Proposition 3.3 there exist  $C_1, C_2, C_3$ ,  $3 \times 3$  matrices such that  $(C_1, C_2, C_3)$  is hypercyclic. Let  $I_3$  be the  $3 \times 3$  identity matrix. Define  $A_1 = \text{diag}(B_1, I_3)$ ,  $A_2 = \text{diag}(B_2, I_3)$ ,  $A_3 = \text{diag}(I_2, C_1)$ ,  $A_4 = \text{diag}(I_2, C_2)$  and  $A_5 = \text{diag}(I_2, C_3)$ . It can easily be shown that  $(A_1, A_2, \dots, A_5)$  is hypercyclic. The details are left to the reader.

## REFERENCES

- [1] F. Bayart, É. Matheron, *Topics in linear dynamics*, Cambridge Tract in maths (to appear).
- [2] N. S. Feldman, *Hypercyclic Tuples of Operators*, Vol. 3, Oberwolfach Rep. 3 (2006), no. 3, 2254-2256. Mini-workshop: Hypercyclicity and Linear Chaos. Abstracts from the mini-workshop held August 13-19, 2006. Organized by Teresa Bermudez, Gilles Godefroy, Karl-G. Grosse-Erdmann and Alfredo Peris. Oberwolfach Reports. Vol. 3, no. 3. Oberwolfach Rep. 3 (2006), no. 3, 2227-2276.
- [3] N. S. Feldman, *Hypercyclic pairs of coanalytic Toeplitz operators*, Integral Equations Operator Theory **58** (2007), 153-173.
- [4] N. S. Feldman, *Hypercyclic tuples of operators & somewhere dense orbits*, preprint.
- [5] K. -G. Grosse-Erdmann, *Universal families and hypercyclic operators*, Bull. Amer. Math. Soc. **36** (1999), 345-381.
- [6] K. -G. Grosse-Erdmann, *Recent developments in hypercyclicity*, RACSAM Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. (2003), 273-286.
- [7] K. -G. Grosse-Erdmann, *Dynamics of linear operators*, preprint.
- [8] L. Kerchy, *Cyclic properties and stability of commuting power bounded operators*, Acta Sci. Math. (Szeged) **71** (2005), 299-312.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CRETE, KNOSSOS AVENUE,  
GR-714 09 HERAKLION, CRETE, GREECE

*E-mail address:* costakis@math.uoc.gr

THE SCHOOL OF SCIENCES, EUROPEAN UNIVERSITY CYPRUS, 6 DIOGENES  
STREET, ENGOMI, P.O.BOX 22006, 1516 NICOSIA, CYPRUS

*E-mail address:* d.hadjiloucas@euc.ac.cy

FAKULTÄT FÜR MATHEMATIK, SFB 701, UNIVERSITÄT BIELEFELD, POST-  
FACH 100131, D-33501 BIELEFELD, GERMANY

*E-mail address:* amanouss@math.uni-bielefeld.de